

District	Total Spots Visited	Total Area Surveyed (Acres)	Fungicides Treatment				% Pest Insects								Beneficials/ spike	% Disease Incidence								
			Spots	%	Area	%	Spots Above ETL				Spots Below ETL					Smut	Brown Rust	Yellow Rust	Intensity			Powdery Mildew	Downy Mildew	Others
							Aphid	ABW	AW	Other	Aphid	ABW	AW	Other					Low	Medium	High			

3. **Smut** incidence (0.65% spots) has been observed in the Punjab in districts Gujranwala, Nankana Sahib, Khanewal, M.Garh, T.T.Singh, Faisalabad and Bhakkar.

4. **Brown Rust** incidence (1.01% spots) has been observed in district Ferozwala (9.30% spots), M.b.Din (3.51% spots), Vehari (2.54% spots), D.G.Khan (4.82% spots), B.Pur (1.16% spots), Jhang (3.61% Spots), Chiniot (14.46 % spots), M.Garh (1.72% spots), T.T. Singh (2.07% spots) and Sahiwal (9.91% spots).

5. **Yellow Rust** incidence (40.46% spots) has been observed in all the district in Punjab.

6. Up till now 21.73% spots have been treated with fungicides covering 25.18% area.

RECOMMENDATIONS

1. Farmers are being advised to spray appropriate weedicides on their crops as last resort.
2. Spray suitable fungicide in case of severe infection of Rust.
3. Fungicide Seed treatments are effective in managing Flag Smut