SUMMARY OF GRAM CROP FOR 2ND WEEK OF DECEMBER ,2021																				
District	Total Spots Visited	Area Surveyed (Acres)	(Acres)	Insect Pest %													Disease Incidence %			
				Above ETL						Below ETL						1	Root	Botrytis		
				Pod Borer	Annıd	_	Cut worm	Army worm	Termite	Pod Borer	Aphid	Grass hopper	Cut worm	Army worm	Termite	Blight	Wilt	Rot	Blight	
Mianwali	10	57	6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	20.0	0.0	0.0	
Khushab	19	522	55	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	21.1	0.0	0.0	0.0	0.0	
Bhakkar	67	327	58	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	14.9	0.0	6.0	0.0	0.0	
Jhang	20	83	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.0	0.0	10.0	0.0	0.0	
Chakwal	7	8.375	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Bahawalnagar	13	138	22	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	15.4	0.0	15.4	0.0	0.0	
Muzaffargarh	5	35	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Layyah	21	234	101	0.0	0.0	0.0	0.0	0.0	0.0	9.5	0.0	0.0	0.0	0.0	23.8	0.0	19.0	0.0	0.0	
Total	162	1404.38	247	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.0	0.0	0.0	0.0	13.6	0.0	8.6	0.0	0.0	

OBSERVATION

- $1.\ Total\ 162\ spots\ of\ Gram\ fields\ comprising\ area\ of\ \ 1404.38\ acres\ in\ 8\ districts\ were\ surveyed\ for\ pests..$
- 2. **Pod Borer** infestation below ETL has been observed on 1.2% spots in districts Layyah (9.5% spots).
- 3. **Termite** infestation below ETL has been observed on 13.6% spots in districts Khushab (21.1% spots), Bhakkar (14.9% spots), Jhang (5.0% spots), Bahawalnagar (15.4%) and Layyah (23.8% spots).
- 4. Total 8.6% spots of **Wilt** attack were observed in district Mianwali, Bhakkar, Jhang, Bahawalnagar and Layyah.

RECOMMENDATIONS.

- 1. Irrigate the field to suppress the infestation of termites.
- 2. Control Pod Borer with the spray of spinosad 240 SC @ 100ml/ acre or emamectin benzoate1.9 EC @ 200ml/acre.
- 3. Uproot and burn the plants affected with disease to check its spread.