





**Pest Situation of Sugarcane
Crop in Punjab
(15TH – 31ST DECEMBER 2022)**

GRAPHICAL PEST SITUATION ON SUGARCANE CROP IN PUNJAB FOR THE 2ND FORTNIGHT OF OCTOBER 2022

A- Insect Pest



B- Disease



**FORTNIGHTLY PEST SCOUTING AND FORECAST REPORT OF SUGARCANE
CROP FOR THE 2nd FORTNIGHT OF DECEMBER 2022**

Pest Situation of Sugarcane Pests								
Sr. No.	Pest Name	%age of spots						Remarks
		Current F.Night 2022		Previous F.Night 2022		Corresponding F.Night of 2021		
		AETL	BETL	AETL	BETL	AETL	BETL	
1	BORER	0.00	3.47	0.00	4.04	0.00	2.94	-
2	PYRILLA	0.00	1.16	0.00	2.69	0.00	2.59	-
3	WHITEFLY	0.00	2.29	0.00	0.00	0.00	0.00	-
4	BLACK BUG	0.00	2.89	0.00	2.02	0.00	0.69	-
-5	MITES	0.00	0.00	0.00	0.00	0.00	0.00	-
6	TOKA	0.00	1.73	0.00	1.35	0.00	0.86	-
7	MEALY BUO	0.00	-	0.00	-	0.00	-	-
8	RED ROT	4.05	-	3.03	-	2.42	-	Increasing
9	WHIP SMUT	2.31	-	1.68	-	0.00	-	Increasing
10	MOSAIC VIRUS	0.00	-	0.00	-	0.52	-	-
11	RUST	0.00	-	0.00	-	0.00	-	-

METEOROLOGICAL DATA OF THE FORTNIGHT

Districts	2022				2021			
	Temperature		R.H%	Rainfall (mm)	Temperature		RH%	Rainfall (mm)
	Max.	Min.			Max.	Min.		
Multan	25.4	7.0	71.9	0.0	19.1	5.6	64.5	0.0
Khanewal	22.2	8.2	65.7	0.0	19.9	6.3	65.5	0.0
Vehari	22.4	9.7	67.4	0.0	19.0	6.4	66.6	0.0
Lodhran	20.8	9.6	68.2	0.0	19.1	6.8	67.6	2.0
Sahiwal	18.0	5.9	92.6	0.0	19.0	6.9	92.0	0.0
Pakpattan	18.8	6.0	90.0	0.0	19.4	6.1	88.0	0.0
Okara	18.5	5.8	92.3	0.0	19.2	6.4	90.0	0.0
Bahawalpur	24.6	10.0	51.1	0.0	33.8	18.1	56.9	0.0
Bahawalnagar	20.1	7.2	83.2	0.0	20.8	7.1	81.3	0.0
R.Y. Khan	26.0	9.0	57.0	0.0	24.4	8.8	64.2	0.0
D.G Khan	23.0	10.4	77.8	0.0	23.5	8.1	61.1	0.0
M. Garh	17.5	5.8	68.1	0.0	31.7	18.1	52.8	0.5
Rajanpur	23.7	10.0	39.8	0.0	22.2	7.1	33.9	2.5
Layyah	21.5	8.5	60.0	0.0	16.8	4.1	80.3	0.5
AVERAGE	21.6	8.1	70.4	0.0	22.0	8.3	68.9	0.4

FORECAST FOR THE NEXT FORTNIGHT

BORERS

This pest flourish best at optimum temperature 35-41C° with relative humidity below 65-70%. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

PYRILLA

This pest flourish best at optimum temperature 29-40C with relative humidity 75-84 . The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

WHITEFLY

This pest flourish best at optimum temperature 29-40C with relative humidity 75-84%. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

BLACK BUG

This pest flourish best at optimum temperature 22.9-40.9C with relative R. humidity 46%. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

TOKA

The optimal temperature and RH conditions for the feeding activities are 28.5-33C and 60-70 C° % respectively. The current weather conditions on overall South Punjab basis is as; maximum

temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

MEALY BUG

This pest flourish best at optimum temperature 29-40C with relative humidity 75-84%. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

TERMITES

The optimal temperature and RH conditions for the feeding activities are 35°-40C and 70-80% respectively . The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will sustain during the next fortnight.

MITES

The optimal temperature and RH conditions for the feeding activities are 30°-36C and 20-40% respectively. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will decrease during the next fortnight.

RUST

This disease flourish best at optimum temperature 12-14C with relative humidity below 80-85%. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%**. It is predicted that the population of this pest will sustain during the next fortnight.

RED ROT

This disease flourish best at optimum temperature 25.5-26.5C with relative humidity 60. The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co,**

minimum 8.1 Co with R. humidity 70.4%. It is predicted that the population of this pest will increase during the next fortnight.

WHIP SMUT

This disease flourish best at optimum temperature 14-35C with relative humidity 55-80%. The current weather conditions on overall South Punjab basis is as; maximum temperature **30.8 Co, minimum 16.8 Co with R. humidity 64.2%.** It is predicted that the population of this pest will increase during the next fortnight.

MOSAIC VIRUS

The current weather conditions on overall South Punjab basis is as; maximum temperature **21.6 Co, minimum 8.1 Co with R. humidity 70.4%.** It is predicted that the population of this pest will sustain during the next fortnight.

RECOMMENDATIONS

BORERS MANAGEMENT

- ❖ Install cards of beneficial insects (*Trichogramma*) from the start of crop.
- ❖ Apply granular insecticide carbofuron 3G@ 12-14 kg/acre to the fields where infestation of sugarcane borers observed above ETL.

PYRILLA MANAGEMENT

- ❖ Promote parasitic insects (*Tetrastichus pyrillae*) against eggs of pyrilla and (*Epiricania melanoleuca*) against nymphs and adults of pyrilla.
- ❖ Cut the leaves 6 inch in length from those fields having parasite eggs and pupae in abundance and shift/hang them in fields where parasites are not found.
- ❖ Apply Granular insecticide Carbofuron 3G@ 12-14 kg/acre to the fields where infestation of sugarcane Pyrilla reaches ETL and parasites are not found. Application of granules may be done till the 6 feet height of canes.

WHITEFLY MANAGEMENT

- ❖ Cut severe infested leaves of whitefly and bury in the soil.
- ❖ Install cards of beneficial insects i.e *Chrysoperla carnea*

BLACK BUG MANAGEMENT

- ❖ Avoid the cane fields from water stress.
- ❖ Apply granular insecticide carbofuron 3G@ 12-14 kg/acre

RUST MANAGEMENT

- ❖ Cultivate resistant varieties.
- ❖ Cut and burn the diseased plants/ Plant pests.

WHIP SMUT MANAGEMENT

- ❖ Cut and burn the diseased plants / Plant pests

ECONOMIC THRESHOLD LEVELS (ETLs) OF SUGARCANE PESTS

INSECT PESTS	ECONOMIC THRESHOLD LEVEL
Borers	10% infested canes.
Pyrilla	3 per leaf.
Whitefly	10 per Leaf
Black bug	10 per sheath.
Toka	3 per sweep
Mites	10 per Leaf
Mealy bug	Only presence
Termites	10% damage
Rodents	5 live burrows per acre
Diseases	Only presence