



GRAPHICAL PEST SITUATION ON RICE CROP IN PUNJAB DURING 3rd WEEK OF JUNE, 2022



PEST SITUATION ON RICE CROP IN PUNJAB DURING 3rd WEEK OF JUNE, 2022

Pest Situation of Rice Pests								
Sr. No.	Pest Name	%Age of spots						Remarks
		Current Week 2022		Previous Week 2022		Corresponding week 2021		
		AETL	BETL	AETL	BETL	AETL	BETL	
1	RICE BORER	0.00	0.00	0.00	0.00	0.00	1.56	-
2	LEAF FOLDER	0.00	0.00	0.00	0.00	0.00	0.26	-
3	WPBH	0.00	0.00	0.00	0.00	0.00	0.00	-
4	BPH	0.00	0.00	0.00	0.00	0.00	0.00	-
5	TOKA	0.00	0.52	0.00	0.00	0.52	2.34	-
6	FOOT ROT	0.75	-	0.00	-	0.00	-	Increasing
7	B.L.B	0.00	-	0.00	-	0.00	-	-
8	B.L.S	0.00	-	0.00	-	0.00	-	-
9	SHEATH BLIGHT	0.00	-	0.00	-	0.78	-	-
10	BLAST	0.00	-	0.00	-	0.52	-	-
NO. OF TOTAL SPOTS VISITED			265					
TOTAL AREA VISITED (Acres)			2364					

Tehsil wise percentage of hot spots of Rice Borer

Nil

Tehsil wise percentage of hot spots of Rice Leaf Folder

Nil

Tehsil wise percentage of hot spots of White-Backed Plant Hoppe

Nil

Tehsil wise percentage of hot spots of Brown Plant Hopper

Nil

Tehsil wise percentage of hot spots of Rice Toka

Nil

Tehsil wise percentage of hot spots of Foot Rot

Sr.	TEHSIL	%AGE	Sr.	TEHSIL	%AGE
1	Wazirabad	33.3	2	Phalia	33.3

Tehsil wise percentage of hot spots of Bacterial Leaf Blight

Nil

Tehsil wise percentage of hot spots of Brown Leaf Spots

Nil

Tehsil wise percentage of hot spots of Sheath Blight

Nil

Tehsil wise percentage of hot spots of Rice Blast
Meteorological data of the current week 2021-2022

METEOROLOGICAL DATA								
Districts	2022				2021			
	Temperature		R.H%	Rainfall (mm)	Temperature		RH%	Rainfall (mm)
	Max.	Min.			Max.	Min.		
Gujranwala	37.8	25.6	55.8	5.0	38.3	27.2	57.1	0.0
Hafizbad	35.0	24.0	53.0	0.0	41.0	29.0	45.0	23.0
Sialkot	36.4	23.9	56.8	6.0	39.9	28.4	54.2	0.0
Narowal	39.7	24.7	57.3	15.0	35.0	21.0	61.9	33.0
Gujrat	39.4	29.1	48.0	62.0	37.0	25.9	36.0	15.0
M.B.Din	37.0	25.0	0.5	15.0	38.0	26.0	46.0	11.0
Lahore	19.2	13.1	24.0	7.0	11.3	8.0	17.8	0.1
Sheikhupura	38.6	26.8	48.0	2.0	39.7	27.5	46.0	14.0
Nankana	42.0	32.3	27.3	4.0	35.7	25.4	37.3	7.0
Kasur	36.3	14.5	46.5	2.5	41.3	14.3	29.3	0.0
Faisalabad	38.5	26.4	60.8	20.4	35.8	26.6	77.4	1.6
Jhang	36.5	25.4	61.8	22.9	25.5	34.6	51.31	
Toba Tek Singh	39.1	26.0	67.0	77.4	37.0	25.0	70.0	6.1
Chiniot	31.7	22.9	69.4	36.0	33.3	26.9	52.0	0.0
Multan	36.7	25.9	72.1	11.1	39.8	29.5	71.3	23.4
Khanewal	34.6	24.9	63.9	5.4	40.1	29.4	50.5	0.0
Vehari	35.7	26.3	61.6	7.0	39.7	28.7	49.6	0.0
Lodhran	38.7	25.3	71.9	20.6	38.7	25.6	74.6	28.0
Sahiwal	39.0	27.2	48.0	14.0	36.0	26.0	57.0	7.0
Pakpattan	40.0	26.5	51.0	12.2	39.0	29.0	51.0	19.0
Okara	39.6	26.2	49.6	9.2	37.0	25.0	48.0	9.5
Bahawalpur	39.1	25.8	59.6	28.0	37.4	26.1	60.1	11.4
Bahawalnagar	41.5	28.1	39.3	5.0	34.6	25.4	60.9	37.0
R.Y.Khan	38.3	25.5	52.5	14.0	41.4	27.8	55.0	3.7
D.G. Khan	44.5	34.5	67.0	8.0	43.0	30.5	56.0	0.0
Muzaffar Garh	33.2	22.3	78.9	15.0	41.5	28.0	41.1	1.8
Rajanpur	43.5	30.2	56.0	7.0	40.1	29.0	55.0	9.0
Layyah	44.0	20.0	50.0	37.7	42.1	27.3	42.4	0.0
Tot/Avg	37.70	25.30	53.48	16.76	37.12	26.18	51.92	9.65

Forecast of Rice Pests:

Borer: This pest flourishes best in warm humid climate with optimum temperature 17-30 °C with relative humidity between 45-80%. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may decrease during the coming week as the temperature remain not favorable for the development of this pest.

Leaf Folder: This pest flourishes best in warm humid climate with optimum temperature 25-30°C. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may decrease during the coming week as the temperature remain not favorable for the development of this pest.

White-backed plant hopper: This pest flourishes best in warm humid climate with optimum temperature 25-29°C with relative humidity between 80-90%. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may decrease during the coming week as the temperature remain not favorable for the development of this pest.

Brown plant hopper: This pest flourishes best in warm humid climate with optimum temperature 28-30°C with relative humidity below 80-90%. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may decrease during the coming week as the temperature remain not favorable for the development of this pest.

Toka: This pest flourishes best in warm humid climate with optimum temperature 24-40°C with relative humidity between 30-80%. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may decrease during the coming week as the temperature remain not favorable for the development of this pest.

Foot rot: High humidity and cloudy weather during heading stage are favorable for the development of foot rot of rice. The fungus have a wide range of temperature for optimum growth which is between 30-35 °C. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may increase during the coming week as the temperature remain favorable for the development of this disease.

Bacterial Leaf Blight: Heavy rain, heavy dew, flooding, deep irrigation water are favorable factors for the development of disease. Temperature for optimum growth is between 25-34 °C with relative humidity above 70%. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may increase during the coming week as the temperature remain favorable for the development of this disease.

Brown Leaf spots: Non-flooded and nutrient deficient soils or soils with accumulation of toxic substances are favorable for the development of disease. Temperature for optimum growth is between 16-36 °C with relative humidity from 86-100%. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may increase during the coming week as the temperature remain favorable for the development of this disease.

Sheath Blight: Crop plants during rainy season are more vulnerable to the disease. Temperature for optimum growth is between 28-32 °C with relative humidity from 85-100%. Keeping in view the

temperature for current week and weather forecast of next week, it is predicted that population of this pest may increase during the coming week as the temperature remain favorable for the development of this disease.

Blast: Intermittent [drizzles](#), cloudy weather, more of rainy days, Low night temperature and longer duration of dew are favorable factors for the development of disease. Keeping in view the temperature for current week and weather forecast of next week, it is predicted that population of this pest may increase during the coming week as the temperature remain favorable for the development of this disease.

RECOMMENDATION

RICE BORER MANAGEMENT

- Handpick and destroy egg masses.
- Install light traps up to September to monitor moth population of stem borers.
- Use balanced Fertilizers (NPK) within 45 days after transplanting of nursery.
- Complete application of nitrogen up to 31st August because due to late application of nitrogenous fertilizer, the plant becomes succulent and dark-green which attracts the insects, and helps in their rapid multiplication along with increasing disease incidence.

BOWN LEAF SPOT MANAGEMENT

- Avoid water stress before maturity.
- Control the disease with one of the following pesticides.

S#	Common Name	Brand Name	Dose / Acre
1	Propineb 70 WP	Gift, Cover, Protest	800 gm
2	Mancozeb 80 WP	Shelter, Dithane-M	800 gm
3	Propiconazole 25 EC	Tilt	80 ml

FOOT ROT MANAGEMENT

- Uproot the diseased plants and destroy them.
- Use Potash 1 Bag within 14 days of transplanting.
- Flooding of Copper Sulphate 1.5-2 Kg/Acre.

BACTERIAL LEAF BLIGHT MANAGEMENT

- Use disease free seeds for next crop.
- Spray copper based fungicides without delay when disease incidence is observed.

PADDY BLAST MANAGEMENT

- For leaf blast, re-flood if field has been drained. Maintain water level at 3-4inches to ensure that soil is covered.
- Avoid late use of nitrogenous fertilizers.
- Control the disease with one of the following fungicides;

S#	Common Name	Brand Name	Dose / Acre
1	Kasugamycin 6% WP	Fork	250 gm
2	Trifloxystrobin+Tebuconazole 75%WP	Nativo	65 gm
3	Azoxystrobin 25 % SC	Primacy	200 ml
4	Difenoconazole 250 EC	Score	125 ml

ECONOMIC THRESHOLD LEVELS OF RICE PESTS

INSECT PESTS	ECONOMIC THRESHOLD LEVELS
Borers (White, Yellow & Pink)	0.5% attack on rice nursery while 8-10 Moth/Trap/Night & 5% dead heart on rice crop.
Toka	3 per net on rice nursery & 5 on rice crop.
Leaf Folder	2 rolled leaves per plant in July-August & 3 rolled leaves per plant in September-October.
Brown Plant Hopper	15 Nymphs or Adults per plant in July-August & 20 Nymphs or Adults per plant in September-October. Or 7-10 Nymphs or Adults per net
White Backed Plant Hopper	15 Nymphs or Adults per plant in July-August & 20 Nymphs or Adults per plant in September-October. Or 7-10 Nymphs or Adults per net
Hispa	1 per plant
Diseases	On appearance