

PEST SITUATION ON COTTON CROP IN PUNJAB DURING 4th WEEK OF July, 2023

Whitefly: On overall Punjab basis 8.16% above ETL spots have been observed during current week as compared to 8.30 % % in last week and 24.55% during the same corresponding period of last year. And 89.39% below ETL spots have been observed during current week as compared to 89.32% in last week and 74.28% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of whitefly

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	MIAN CHANNU	46.2	29	KHAN PUR	7.1
2	BHAKKAR	40.0	30	TAUNSA	6.7
3	RAJAN PUR	33.3	31	ARIF WALA	6.5
4	BAHAWAL PUR	24.0	32	HASILPUR	6.3
5	KHANEWAL	23.7	33	CHISHTIAN	6.1
6	CHICHA WATNI	22.4	34	RENALA KHURD	5.9
7	SILANWALI	18.8	35	CHOBARA	5.7
8	BURE WALA	17.8	36	SUMUNDRI	5.6
9	ROJHAN	17.1	37	LAYYAH	5.0
10	T.T.SINGH	16.9	38	BHERA	4.7
11	VEHARI	16.3	39	JAHANIAN	4.7
12	KABIR WALA	15.7	40	MINCHAN ABAD	4.7
13	JAM PUR	15.0	41	MUZUFAR GARH	4.3
14	PIR MAHAL	13.6	42	KAROR L.E.	4.3
15	QUIDABAD	12.5	43	BAHAWAL NAGAR	4.2
16	KOT CHUTTA	12.1	44	PIPLAN	4.0
17	KEHROR PACCA	11.1	45	OKARA	3.8
18	JHANG	11.1	46	PAK PATTAN	3.7
19	BHALWAL	10.0	47	MULTAN	3.6
20	MAILSI	9.8	48	LIAQUAT PUR	3.6
21	ISA KHEL	9.5	49	D.G.KHAN	3.1
22	R.Y.KHAN	9.4	50	JARANWALA	2.5
23	KOT ADDU	8.6	51	TANDLIANWALA	2.5
24	ALI PUR	8.5	52	DUYAPUR	2.3
25	KHUSHAB	8.0	53	FORT ABBAS	2.1
26	SAHIWAL	7.9	54	GOJRA	1.6
27	SADIQ ABAD	7.7	55	JATOI	1.4
28	HAROON ABAD	7.1			

Jassid: On overall Punjab basis 15.21% above ETL spots have been observed as compared to 15.83% in last week and same week of last year 19.69% and 83.28% below ETL spots have been observed during current week as compared to 82.95% in last week and 77.79% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of Jassid

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	SAHIWAL	66.7	34	A.P.EAST	16.0
2	JALAL PUR P.W	47.9	35	MIANWALI	15.8
3	BAHAWAL PUR	46.0	36	T.T.SINGH	15.7
4	SHUJABAD	44.6	37	ALI PUR	15.5
5	KOTMOMIN	33.3	38	HAROON ABAD	14.3
6	MULTAN	32.1	39	VEHARI	14.0
7	SILANWALI	31.3	40	JHANG	13.9
8	YAZMAN	27.1	41	PIR MAHAL	13.6
9	DUYAPUR	25.0	42	SADIQ ABAD	13.5

10	DARYA KHAN	25.0	43	QUIDABAD	12.5
11	LODHRAN	23.8	44	BHERA	12.5
12	TAUNSA	23.3	45	D.G.KHAN	12.5
13	FORT ABBAS	22.9	46	BHALWAL	10.0
14	CHISHTIAN	22.4	47	KABIR WALA	9.8
15	CHOBARA	21.4	48	ROJHAN	9.8
16	KAROR L.E.	21.4	49	KEHROR PACCA	8.9
17	KOT CHUTTA	21.2	50	JATOI	8.6
18	BHAKKAR	20.0	51	SHORKOT	8.5
19	KOT ADDU	20.0	52	KHANEWAL	8.5
20	MUZUFAR GARH	20.0	53	SAHIWAL	7.9
21	SARGODHA	20.0	54	JARANWALA	7.4
22	ISA KHEL	19.0	55	TANDLIANWALA	7.4
23	R.Y.KHAN	18.9	56	GOJRA	6.5
24	HASILPUR	18.8	57	MIAN CHANNU	5.8
25	BURE WALA	17.8	58	A.P SIAL	5.2
26	MAILSI	17.1	59	JAHANIAN	4.7
27	BAHAWAL NAGAR	16.7	60	SUMUNDRI	4.2
28	MINCHAN ABAD	16.3	61	OKARA	3.8
29	KHAN PUR	16.1	62	PAK PATTAN	3.7
30	LIAQUAT PUR	16.1	63	JAM PUR	3.3
31	KHUSHAB	16.0	64	ARIF WALA	3.2
32	LAYYAH	16.0	65	CHICHA WATNI	2.6
33	PIPLAN	16.0	66	FAISALABAD	2.6

Thrips: On overall Punjab basis 7.09% above ETL spots have been observed as compared to 7.47% in last week and same week of last year i.e. 2.48%. And 90.25% below ETL spots have been observed during current week as compared to 91.33% in last week and 93.52% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of Thrips

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	BAHAWAL PUR	32.0	22	KHUSHAB	8.0
2	SHUJABAD	26.8	23	D.G.KHAN	7.8
3	FORT ABBAS	25.0	24	HAROON ABAD	7.1
4	MUZUFAR GARH	22.9	25	VEHARI	7.0
5	MULTAN	21.4	26	SILANWALI	6.3
6	JATOI	21.4	27	LODHRAN	6.3
7	R.Y.KHAN	17.0	28	KAROR L.E.	5.7
8	CHISHTIAN	16.3	29	KOT ADDU	5.7
9	LIAQUAT PUR	16.1	30	ISA KHEL	4.8
10	TAUNSA	15.0	31	JAHANIAN	4.7
11	JALAL PUR P.W	14.6	32	KOT CHUTTA	4.5
12	ALI PUR	14.1	33	KEHROR PACCA	4.4
13	MINCHAN ABAD	14.0	34	HASILPUR	4.2
14	DUYAPUR	13.6	35	A.P.EAST	4.0
15	SADIQ ABAD	11.5	36	PAK PATTAN	3.7
16	KHAN PUR	10.7	37	BHERA	3.1
17	MIANWALI	10.5	38	MAILSI	2.4
18	BAHAWAL NAGAR	10.4	39	GOJRA	1.6
19	LAYYAH	10.0	40	CHOBARA	1.4
20	BURE WALA	8.9	41	T.T.SINGH	1.2
21	YAZMAN	8.3			

Mealy Bug: On overall Punjab basis 2.21% spots (all considered above ETL) of Mealy Bug have been observed as compared to 2.16% in last week and 5.61% spots during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of mealy bug

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	R.Y.KHAN	17.0	15	ROJHAN	2.4
2	JALAL PUR P.W	14.6	16	VEHARI	2.3
3	RAJAN PUR	13.9	17	DUYAPUR	2.3
4	ALI PUR	11.3	18	FORT ABBAS	2.1
5	OKARA	7.7	19	SADIQ ABAD	1.9
6	KOT CHUTTA	7.6	20	MIAN CHANNU	1.9
7	JATOI	7.1	21	D.G.KHAN	1.6
8	YAZMAN	6.3	22	MUZUFAR GARH	1.4
9	SHUJABAD	5.4	23	KOT ADDU	1.4
10	KHAN PUR	5.4	24	CHOBARA	1.4
11	A.P.EAST	4.0	25	JARANWALA	1.2
12	JAM PUR	3.3	26	TANDLIANWALA	1.2
13	LAYYAH	3.0	27	T.T.SINGH	1.2
14	KAROR L.E.	2.9			

Mites: On overall Punjab basis 0.00% above ETL spots has been observed during current week as compared to 0.00% in last week and 0.00% during the same corresponding period of last year. And 5.64% below ETL spots have been observed during current week as compared to 5.28% in last week and 0.87% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of mites

No hot spot observed this week

Dusky Cotton Bug: On overall Punjab basis 0.40% above ETL spots have been observed during current week as compared to 0.24% in last week and 0.49% during the same corresponding period of last year. And 23.87% below ETL spots have been observed during current week as compared to 17.53% in last week and 26.77% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of dusky cotton bug

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	SHUJABAD	10.7	4	DUYAPUR	2.3
2	ALI PUR	4.2	5	JATOI	1.4
3	JALAL PUR P.W	4.2			

Armyworm: On overall Punjab basis 0.31% spots (all considered above ETL) of Armyworm have been observed as compared to 0.21% in last week and 0.60% spots during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of army worm

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	HAROON ABAD	7.1	4	CHISHTIAN	2.0
2	FORT ABBAS	6.3	5	BAHAWAL PUR	2.0
3	MAILSI	4.9	6	BHERA	1.6

Pink Boll Worm: On overall Punjab basis 0.95% above ETL spots have been observed as compared to 0.43% in last week and 0.11% same corresponding period of the last year. And 15.55% below ETL spots have been observed as compared to 11.69% in last week and 5.46% during same corresponding period of last year.

Tehsil wise percentage of hot spots of pink boll worm

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	HAROO ABAD	7.1	10	D.G.KHAN	3.1
2	ALI PUR	7.0	11	JATOI	2.9
3	R.Y.KHAN	5.7	12	MAILSI	2.4
4	RAJAN PUR	5.6	13	VEHARI	2.3
5	KEHROR PACCA	4.4	14	FORT ABBAS	2.1
6	JALAL PUR P.W	4.2	15	CHISHTIAN	2.0
7	OKARA	3.8	16	PIR MAHAL	1.7
8	ROJHAN	3.7	17	TAUNSA	1.7
9	ARIF WALA	3.2	18	MUZUFAR GARH	1.4

American Boll Worm: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and 0.00% during same corresponding period of last year. And 0.52% below ETL spots have been observed as compared to 0.53% in last week and 0.26% during same corresponding period of last year.

Tehsil wise percentage of hot spots of american boll worm

No hot spot find yet.

Spotted Boll Worm: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and 0.04% during same corresponding period of last year & 0.28% below ETL spots have been observed as compared to 0.21% in last week and 0.15% during same corresponding period of last year.

Tehsil wise percentage of hot spots of spotted boll worm

No hot spot find yet.

CLCuV: On overall Punjab basis 0.61% spots (all considered above ETL) of CLCV have been observed as compared to 0.43% in last week and 31.66% during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of CLCuV

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	BHALWAL	10.0	8	JALAL PUR P.W	2.1
2	SHUJABAD	7.1	9	BAHAWAL PUR	2.0
3	VEHARI	4.7	10	KHAN PUR	1.7
4	MUZUFAR GARH	4.3	11	BHERA	1.6
5	LIAQUAT PUR	3.6	12	JATOI	1.4
6	MAILSI	2.4	13	LODHRAN	1.3
7	KEHROR PACCA	2.2			

FORECAST FOR THE NEXT WEEK

Meteorological data of the current week 2022-2023

Districts	2023				2022			
	Temperature °C		R.H%	Rainfall (mm)	Temperature °C		RH%	Rainfall (mm)
	Max.	Min.			Max.	Min.		
Multan	36.7	27.6	62.8	3.8	34.9	26.4	77.4	8.8
Khanewal	34.7	29.1	68.3	1.1	31.9	26.9	78.5	3.4
Vehari	34.4	27.7	76.6	14.4	33.4	26.9	83.1	8.1
Lodhran	36.0	25.5	73.1	11.1	31.7	24.4	80.4	13.4
Sahiwal	34.7	26.1	82.0	145.0	35.0	25.0	88.0	46.8
Pakpattan	34.5	27.4	78.6	40.0	34.6	26.2	86.0	35.5
Okara	34.3	24.4	82.5	218.2	36.2	26.4	85.5	35.3
Bahawalpur	36.1	26.7	84.4	77.5	35.6	26.0	82.1	127.7
Bahawalnagar	35.7	26.4	73.0	9.2	36.8	28.6	68.7	10.0
R.Y.Khan	34.9	27.6	71.4	24.2	31.9	26.3	75.5	31.4
D.G. Khan	35.3	28.0	71.2	8.0	33.1	26.3	89.1	15.0
Muzaffar Garh	35.3	27.5	74.9	24.0	33.2	26.9	79.3	20.0
Rajanpur	35.3	27.7	80.5	12.0	32.7	26.0	85.5	34.0
Layyah	37.0	21.0	70.0	10.0	34.7	20.0	89.0	81.3
TOT/AVG	35.35	26.62	75.09	599	33.97	25.87	81.99	471

Weather forecast for next 7 day in cotton zone

Division	Dated	1/8	2/8	3/8	4/8	5/8	6/8	7/8
Multan	Max.Temp.	41	40	41	42	40	39	40
	Min.Temp.	31	32	32	32	32	30	31
	Humidity %	36	38	34	35	38	41	37
Sahiwal	Max.Temp.	38	38	39	38	37	38	38
	Min.Temp.	30	30	30	30	30	30	31
	Humidity %	46	47	45	48	50	44	44
B. pur	Max.Temp.	40	39	39	40	40	39	37
	Min.Temp.	31	31	31	31	31	31	30
	Humidity %	42	40	39	39	37	40	46
D.G. Khan	Max.Temp.	41	41	41	41	42	41	40
	Min.Temp.	30	32	31	32	32	32	31
	Humidity %	33	37	37	36	34	38	41
Average Cotton Region	Max.Temp.	39.64						
	Min.Temp.	30.96						
	Humidity %	40.07						

Source: timeanddate.com

Summary of weather forecast

Overall weather for seven days in cotton zone during next week will remain hot, sunny & shiny.

Forecast of Sucking Pests:

Whitefly: This pest flourishes best in hot dry climate with optimum temperature 35-42 °C with relative humidity below 50%. Maximum lethal temperature for this pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%. Keeping in view above all, it is predicted that population of this pest will increase during the coming week.

Jassid: This pest flourishes best in hot and humid climate with optimum temperature 35-40 °C with relative humidity above 70%. Maximum lethal temperature for this pest is 45-51 °C .The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%.Keeping in view above all, it is predicted that population of this pest will increase during the coming week.

Thrips: This pest flourishes best in hot dry climate with optimum temperature 35-40 °C with relative humidity below 60%. Maximum lethal temperature for this pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%.In view above all, it is predicted that population of this pest will decrease during the coming week.

Mealy Bug: This pest flourishes best in moderate climate with optimum temperature 35-40 °C with relative humidity above 60-80%. Maximum lethal temperature for this pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%.Keeping in view above all, it is

predicted that population of this pest will increase as such on cotton crop during the coming week.

Mites: This pest flourishes best in dry climate with optimum temperature 28-40 °C with relative humidity above 40-50%. Maximum lethal temperature for this pest is 45 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%. Keeping in view above all, it is predicted that population of this pest will sustain as such during the coming week.


Dusky Cotton Bug: This pest flourishes best in moderate climate with optimum temperature 28-40 °C with relative humidity above 50-80%. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%. In view above all, it is predicted that population of this pest may sustain during the coming week.

Pink Bollworm: This pest flourishes best at moderate climate with optimum temperature 32-40 °C with relative humidity above 60-80%. Maximum lethal temperature for this pest is 45-53 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%. It is predicted that population of this pest will increase on cotton during the coming week.

Moth Catches in Pheromone Traps during 4th Week of July, 2023

Moth Catches in Pheromone Traps During 4th Week of July, 2023			
District	Traps Installed	Moth catches/Trap/week	Av. Moth catches/Trap/night
Multan	6	41	0.98
Khanewal	8	57	1.02
Vehari	6	49	1.17
Lodhran	6	43	1.02
Sahiwal	4	16	0.57
Pakpattan	4	15	0.54
Okara	4	16	0.57
Bahawalpur	8	47	0.84
Bahawalnagar	10	56	0.80
R.Y.Khan	8	46	0.82
D.G.Khan	6	20	0.48
Rajanpur	6	22	0.52
M.Garh	8	27	0.48
Layyah	6	20	0.48
Bhakkar	8	30	0.54
Mianwali	6	25	0.60
Faisalabad	6	77	1.83
TT. Singh	6	06	0.14
Jhang	4	22	0.79

Note. 3 moth catches on 3 consecutive nights in a trap is considered as ETL



American Bollworm: This pest flourishes best at dry & moderate climate with optimum temperature 32-40 °C with relative humidity above 50-70%. Maximum lethal temperature for this pest is 45-53 °C .The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity

40.07%.It is predicted that population of this pest will remain sustain on cotton crop during the coming week.

Moth catches on light traps during 4rth week of July, 2023

Moth Catches on Light Traps/Trap/night During 4rth Week of July, 2023				
District	Traps Installed	A.B.W	S.B.W	Armyworm
Multan	3	1.3	0.0	2.6
Khanewal	4	2.1	0.0	6.0
Vehari	3	1.1	0.0	2.3
Lodhran	3	3.4	0.0	4.1
Sahiwal	2	0.4	0.0	1.1
Pakpattan	2	0.4	0.0	1.0
Okara	3	0.6	0.0	1.4
Bahawalpur	4	1.7	1.0	1.6
Bahawalnagar	5	1.0	0.7	2.0
R.Y.Khan	4	0.9	0.9	1.1
D.G.Khan	3	0.3	0.1	1.0
M.Garh	4	0.0	0.0	1.4
Rajan Pur	3	0.6	0.4	1.3
Layyah	3	0.1	0.0	0.6
Note. 3 moth catches of a pest on 3 consecutive nights in a trap is considered as ETL				

Spotted Bollworm: This pest flourishes best at mild & humid climate with optimum temperature 30-40 °C with relative humidity above 60-100%. Maximum lethal temperature for this pest is 45-53 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C with humidity 40.07%.Due to cultivation of more BT varieties, it is predicted that population of this pest will remain as such on cotton crop in the coming week.

Armyworm: This pest flourishes best in mild & humid climate with optimum temperature 30-40 °C with relative humidity below 70-100%. The current weather conditions on overall Punjab basis (temp. Max. 35.35 °C, Min. 26.62 °C with R. humidity 75.09%) & forecast for the next week temp. Max. 39.64 °C, Min. 30.96°C

with humidity 40.07%. Keeping in view above all, it is predicted that population of this pest will sustain during the coming week.

CLCuV: The incidence of this disease is expected to sustain as such on cotton crop during the next week.

RECOMMENDATIONS

WHITEFLY MANAGEMENT

1. Conduct regular pest scouting of Whitefly.
2. Control the pest without delay when population reaches ETL i.e. nymphs or adults or both 5 per leaf.
3. Control weeds in and outside the field especially near water channels and fallow lands.

Alternate host plants of whitefly

Crops	Vegetables	Orchards	Weeds/ Ornamentals	Trees
Sunflower	Okra	Citrus	Gardenia	Shisham
Tobacco	Brinjal	Litchi	Mako	Shareen
	Cucurbits	Pomegranate	Maina	
	Tomato	Ber (<i>Zizyphus</i>)	Karund	
	Cabbage	Guava	Lehli	
	Cauliflower	Mulberry		
	Peas	Papaya		
	Potato			
	Onion			
	Spinach			

JASSID MANAGEMENT

1. Eradicate weeds.
2. Conduct Pest Scouting at least twice a week.

3. Use *Chrysoperla* Cards @ 80-90 cards per acre having 20-25 eggs on each card which are available at AARI Faisalabad, Sahiwal, Okara and Vehari.
(Repeat this process after every two weeks).

THRIPS MANAGEMENT

1. Conduct Pest Scouting regularly at least twice a week.
2. Avoid intercropping of cotton crop with vegetables.
3. Control weeds in and outside the field.

MEALY BUG MANAGEMENT

1. Visit the fields daily to detect the infestation for effective management of mealy bug well in time.
2. Keep field and water courses etc. free from weeds especially Itsit & Hazaardani, kangji booti, aksin etc.
3. To avoid its shifting from one place to another, keep the horticultural nurseries free from its infestation. Spray such infested nurseries before shifting the plants.
4. Avoid water stress to the plants.
5. Avoid repeated visits of workers from infested fields to healthy fields.
6. Apply the localized chemical spray on infected along with its 2 meter surrounding area.

Alternate host plants of cotton mealy bug

Crops	Vegetables	Ornamentals	Weeds	Orchards
Sunflower	Okra	China Rose	Hazar Dani	Citrus
Tobacco	Brinjal	Huddle	Amarantus	Mulberry
Jantar	Tomato	Cotton Rose	Bhakra	Ficus
Sesame	Chillies	Gulchain	Mako	Ber
	Pumpkin	Lantana	Sueda	
		Din Ka Raja	Itsit	
		Rat Ki Rani	Karund	
		Anthorium	Aksen	

		Gul-e-Daudi	Bathu	
		Gainda	Puth Kanda	
			Kanghi	

MANAGEMENT OF DUSKY COTTON BUG

1. Keep the special observation/ pest scouting when the boll opening starts.
2. Use of soft contact poison on the above ETL.

MANAGEMENT OF ARMY WORM

1. Keep the fields clean from weeds especially Its it which is a preferred host of armyworm.
2. Observe the field on regular basis and under close observation to assess its attack and treat the patch at earliest if necessary.
3. Avoid sowing of Jantar near cotton fields.
4. Regular pest scouting at least twice a week.
5. Hand picking and destruction of egg masses at initial stage as initially its attack starts in patches and is clearly visible.

CLCuV MANAGEMENT

1. Use CLCuV tolerant varieties for sowing.
2. Keep Cotton fields free from all kinds of weeds.
3. Eradicate other alternate host plants of CLCuV and dispose them off carefully.
4. Thinning out and destruction of Virus affected plants.
5. Remove CLCuV affected plants in less than 60 DAS (Days after sowing) crop.
6. Hoeing after each irrigation or rain at watar condition till canopies permit.

7. Apply post emergence Glyphosate for both broad and narrow leaved weeds and Glyphosate for grassy weeds with shield and use Flat Fan Nozzle, if needed.
8. Irrigate the fields when needed keeping in view weather conditions e.g. temperature, rainfall, soil water holding capacity and plant need.
9. Balanced use of Fertilizer i.e. N.P.K. along with foliar growth promoters if necessary.
10. Keep transmission vector (whitefly) at the low ebb

Alternate host plants of CLCuV vector (whitefly)

Crops	Vegetables	Ornamentals	Weeds
Sunflower	Okra	Gurhal	Leh
Melon	Brinjal	Chambeli	Lehli
Tobacco	Chillies		Mako
	Tomato		Maina
	Potato		Karund/Bathu
	Cucumber		Gardenia
			Hazardani
			Rattanjot
			Sun Kukra

Economic threshold levels of cotton pests

Insect pests	Economic threshold levels
Jassid	1 Adult or Nymph per leaf
Whitefly	5 Adults or Nymphs or both per Leaf
Thrips	8-10 Adults or Nymphs per Leaf
Mealy bug	0 (on appearance)
Mites	On damage appearance
Spotted Boll worm	3 Larvae/ 25 plants.
Pink Boll worm	5 Larvae /100 bolls.
American boll worm	5 brown eggs or 3 small larvae or both 5 per 25 plants on non Bt varieties. 2 larvae of 2 nd instar per 25 plants on Bt varieties.
Army worm	0 (On appearance)
Aphid	Spray on visible damage on top terminals.
Dusky Cotton bug	10 per leaf/bud

