PEST SITUATION ON COTTON CROP IN PUNJAB DURING 2nd WEEK OF OCTOBER, 2023

Whitefly: On overall Punjab basis 10.09% above ETL spots have been observed during current week as compared to 13.25% % in last week and 3.75% during the same corresponding period of last year. And 84.24% below ETL spots have been observed during current week as compared to 84.31% in last week and 82.03% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of whitefly

| S. NO. | TEHSIL | %AGE | S.NO. | TEHSIL | %AGE |
|--------|---------------|------|-------|---------------|------|
| 1 | SADIQ ABAD | 40.9 | 18 | MUZUFAR GARH | 11.9 |
| 2 | R.Y.KHAN | 36.1 | 19 | JALAL PUR P.W | 11.1 |
| 3 | A.P.EAST | 34.0 | 20 | KOT ADDU | 11.1 |
| 4 | LIAQUAT PUR | 32.8 | 21 | ALI PUR | 11.0 |
| 5 | HASILPUR | 29.3 | 22 | SHUJABAD | 10.0 |
| 6 | KHAN PUR | 29.2 | 23 | BURE WALA | 10.0 |
| 7 | YAZMAN | 27.0 | 24 | VEHARI | 8.3 |
| 8 | BAHAWAL PUR | 26.6 | 25 | KEHROR PACCA | 8.3 |
| 9 | CHISHTIAN | 22.2 | 26 | LODHRAN | 5.5 |
| 10 | FORT ABBAS | 20.6 | 27 | KAROR L.E. | 5.4 |
| 11 | BAHAWAL NAGAR | 20.0 | 28 | TAUNSA | 4.2 |
| 12 | MINCHAN ABAD | 20.0 | 29 | LAYYAH | 4.2 |
| 13 | HAROON ABAD | 19.0 | 30 | DARYA KHAN | 3.1 |
| 14 | JATOI | 13.6 | 31 | KALLOR KOT | 2.3 |
| 15 | MULTAN | 13.5 | 32 | DUYAPUR | 2.1 |
| 16 | CHOBARA | 12.5 | 33 | JAHANIAN | 1.9 |
| 17 | MAILSI | 12.1 | | | |

Jassid: On overall Punjab basis 0.31% above ETL spots have been observed as compared to 0.48% in last week and same week of last year 2.27% and 80.05% below ETL spots have been observed during current week as compared to 94.03% in last week and 75.84% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of Jassid

| S. NO. | TEHSIL | %AGE | S.NO. | TEHSIL | %AGE |
|--------|------------|------|-------|-------------|------|
| 1 | QASUR | 33.3 | 5 | T.T.SINGH | 2.9 |
| 2 | CHUNIAN | 25.0 | 6 | KALLOR KOT | 2.3 |
| 3 | SAHIWAL | 3.6 | 7 | KABIR WALA | 2.1 |
| 4 | DARYA KHAN | 3.1 | 8 | BAHAWAL PUR | 1.6 |

Thrips: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and same week of last year i.e. 0.00%. And 56.27% below ETL spots have been observed during current week as compared to 73.44% in last week and 53.28% during the same corresponding period of last year.

Mealy Bug: On overall Punjab basis 10.99% spots (all considered above ETL) of Mealy Bug have been observed as compared to 11.97% in last week and 4.59% spots during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of mealy bug

| S. NO. | TEHSIL | %AGE | S.NO. | TEHSIL | %AGE |
|--------|---------------|------|-------|--------------|------|
| 1 | BAHAWAL PUR | 56.3 | 20 | TAUNSA | 12.5 |
| 2 | R.Y.KHAN | 37.5 | 21 | LODHRAN | 11.0 |
| 3 | KHAN PUR | 34.7 | 22 | KOT CHUTTA | 10.4 |
| 4 | SADIQ ABAD | 30.3 | 23 | MINCHAN ABAD | 10.0 |
| 5 | HASILPUR | 29.3 | 24 | JATOI | 9.1 |
| 6 | YAZMAN | 27.0 | 25 | KAROR L.E. | 8.9 |
| 7 | LIAQUAT PUR | 26.6 | 26 | RAJAN PUR | 8.7 |
| 8 | A.P.EAST | 24.0 | 27 | KABIR WALA | 8.3 |
| 9 | MULTAN | 23.1 | 28 | CHOBARA | 8.3 |
| 10 | ROJHAN | 22.2 | 29 | KEHROR PACCA | 6.3 |
| 11 | BAHAWAL NAGAR | 20.0 | 30 | DUYAPUR | 6.3 |
| 12 | SHUJABAD | 20.0 | 31 | D.G.KHAN | 6.3 |
| 13 | CHISHTIAN | 18.5 | 32 | LAYYAH | 5.6 |
| 14 | JALAL PUR P.W | 16.7 | 33 | ALI PUR | 4.0 |
| 15 | VEHARI | 16.7 | 34 | JAM PUR | 3.7 |
| 16 | FORT ABBAS | 15.9 | 35 | MIAN CHANNU | 3.6 |
| 17 | MAILSI | 15.5 | 36 | BHAKKAR | 3.3 |
| 18 | BURE WALA | 15.0 | 37 | GOJRA | 2.4 |
| 19 | HAROON ABAD | 14.3 | 38 | MUZUFAR GARH | 1.7 |

Mites: On overall Punjab basis 0.00% above ETL spots has been observed during current week as compared to 0.00% in last week and 0.00% during the same corresponding period of last year. And 2.27% below ETL spots have been observed during current week as compared to 2.01% in last week and 1.85% during the same corresponding period of last year.

Dusky Cotton Bug: On overall Punjab basis 8.24% above ETL spots have been observed during current week as compared to 6.45% in last week and 9.93% during the same corresponding period of last year. And 60.11% below ETL spots have been observed during current week as compared to 60.64% in last week and 66.75% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of dusky cotton bug

| S. NO. | TEHSIL | %AGE | S.NO. | TEHSIL | %AGE |
|--------|---------------|------|-------|-------------|------|
| 1 | BAHAWAL PUR | 92.2 | 15 | T.T.SINGH | 8.8 |
| 2 | HASILPUR | 53.7 | 16 | MANKERA | 8.0 |
| 3 | A.P.EAST | 44.0 | 17 | PIPLAN | 6.3 |
| 4 | YAZMAN | 36.5 | 18 | VEHARI | 5.6 |
| 5 | SHUJABAD | 28.0 | 19 | LODHRAN | 5.5 |
| 6 | MULTAN | 26.9 | 20 | MIAN CHANNU | 5.5 |
| 7 | JALAL PUR P.W | 22.2 | 21 | KALLOR KOT | 4.5 |
| 8 | HAROON ABAD | 22.2 | 22 | KHANEWAL | 4.1 |
| 9 | FORT ABBAS | 20.6 | 23 | MAILSI | 3.4 |
| 10 | MIANWALI | 14.3 | 24 | BURE WALA | 2.5 |
| 11 | CHISHTIAN | 11.1 | 25 | DUYAPUR | 2.1 |
| 12 | JATOI | 10.6 | 26 | ALI PUR | 2.0 |
| 13 | ISA KHEL | 10.0 | 27 | JAHANIAN | 1.9 |
| 14 | PIR MAHAL | 9.7 | 28 | SUMUNDRI | 1.1 |

Armyworm: On overall Punjab basis 0.00% spots (all considered above ETL) of Armyworm have been observed as compared to 0.11% in last week and 0.21% spots during the same corresponding period of the last year.

Pink Boll Worm: On overall Punjab basis 15.48% above ETL spots have been observed as compared to 15.15% in last week and 18.60% same corresponding period of the last year. And 43.53% below ETL spots have been observed as compared to 43.16% in last week and 42.97% during same corresponding period of last year.

Tehsil wise percentage of hot spots of pink boll worm

| S. NO. | TEHSIL | %AGE | S.NO. | TEHSIL | %AGE |
|--------|---------------|------|-------|--------------|------|
| 1 | A.P.EAST | 52.0 | 26 | ALI PUR | 15.0 |
| 2 | BAHAWAL PUR | 50.0 | 27 | DUYAPUR | 12.5 |
| 3 | SHUJABAD | 34.0 | 28 | TAUNSA | 12.5 |
| 4 | R.Y.KHAN | 33.3 | 29 | KOT CHUTTA | 12.5 |
| 5 | KHAN PUR | 33.3 | 30 | KABIR WALA | 12.5 |
| 6 | MULTAN | 32.7 | 31 | KEHROR PACCA | 12.5 |
| 7 | JALAL PUR P.W | 31.5 | 32 | D.G.KHAN | 12.5 |
| 8 | HASILPUR | 29.3 | 33 | KOT ADDU | 12.5 |
| 9 | SADIQ ABAD | 28.8 | 34 | KAROR L.E. | 10.7 |
| 10 | LIAQUAT PUR | 28.1 | 35 | MIANWALI | 9.5 |
| 11 | JATOI | 24.2 | 36 | T.T.SINGH | 8.8 |
| 12 | YAZMAN | 23.8 | 37 | CHOBARA | 8.3 |
| 13 | KHANEWAL | 22.4 | 38 | MANKERA | 8.0 |
| 14 | FORT ABBAS | 20.6 | 39 | LAYYAH | 6.9 |
| 15 | CHISHTIAN | 20.4 | 40 | BHAKKAR | 6.7 |
| 16 | MIAN CHANNU | 20.0 | 41 | PIPLAN | 6.3 |
| 17 | BAHAWAL NAGAR | 20.0 | 42 | KALLOR KOT | 4.5 |
| 18 | MINCHAN ABAD | 20.0 | 43 | ROJHAN | 3.7 |
| 19 | LODHRAN | 19.2 | 44 | JAM PUR | 3.7 |
| 20 | HAROON ABAD | 19.0 | 45 | PIR MAHAL | 3.2 |
| 21 | MAILSI | 19.0 | 46 | DARYA KHAN | 3.1 |
| 22 | BURE WALA | 17.5 | 47 | PAK PATTAN | 2.1 |
| 23 | VEHARI | 16.7 | 48 | SAHIWAL | 1.8 |
| 24 | MUZUFAR GARH | 15.3 | 49 | SUMUNDRI | 1.1 |
| 25 | JAHANIAN | 15.1 | 50 | TANDLIANWALA | 1.1 |

American Boll Worm: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and 0.00% during same corresponding period of last year. And 0.24% below ETL spots have been observed as compared to 0.19% in last week and 1.14% during same corresponding period of last year.

Spotted Boll Worm: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and 0.00% during same corresponding period of last year & 0.14% below ETL spots have been observed as compared to 0.24% in last week and 0.34% during same corresponding period of last year.

CLCuV: On overall Punjab basis 2.44% spots (all considered above ETL) of CLCV have been observed as compared to 3.29% in last week and 12.54% during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of CLCuV

| S. NO. | TEHSIL | %AGE | S.NO. | TEHSIL | %AGE |
|--------|-------------|------|----------------|--------------|------|
| 1 | KABIR WALA | 27.1 | 11 | PIPLAN | 6.3 |
| 2 | VEHARI | 16.7 | 12 | KEHROR PACCA | 4.2 |
| 3 | MIAN CHANNU | 14.5 | 13 | KHANEWAL | 4.1 |
| 4 | MIANWALI | 14.3 | 14 | ALI PUR | 4.0 |
| 5 | BURE WALA | 12.5 | 15 | BHAKKAR | 3.3 |
| 6 | JATOI | 12.1 | 16 | T.T.SINGH | 2.9 |
| 7 | MAILSI | 12.1 | 17 | LODHRAN | 2.7 |
| 8 | ISA KHEL | 10.0 | 18 JAHANIAN | | 1.9 |
| 9 | MANKERA | 8.0 | 19 BAHAWAL PUR | | 1.6 |
| 10 | KALLOR KOT | 6.8 | | | |

FORECAST FOR THE NEXT WEEK

Meteorological data of the current week 2022-2023

| | | 2023 | | | 2022 | | | |
|---------------|----------------|-------|----------|------|---------|----------------|-------|----------|
| Districts | Temperature °C | | Rainfall | | Tempera | Temperature °C | | Rainfall |
| | Max. | Min. | R.H% | (mm) | Max. | Min. | RH% | (mm) |
| Multan | 37.0 | 22.0 | 65.0 | 0.0 | 35.0 | 21.6 | 66.3 | 0.0 |
| Khanewal | 35.7 | 23.6 | 53.9 | 0.0 | 35.3 | 22.4 | 65.6 | 0.0 |
| Vehari | 35.6 | 23.6 | 50.0 | 0.0 | 34.1 | 20.9 | 70.2 | 0.0 |
| Lodhran | 35.3 | 23.9 | 64.8 | 0.0 | 33.6 | 22.8 | 71.0 | 0.0 |
| Sahiwal | 35.5 | 21.0 | 63.0 | 0.0 | 31.4 | 19.0 | 67.5 | 0.0 |
| Pakpattan | 32.2 | 24.6 | 36.0 | 0.0 | 40.8 | 20.8 | 54.4 | 0.0 |
| Okara | 33.0 | 22.5 | 59.7 | 0.0 | 33.7 | 23.3 | 60.2 | 0.0 |
| Bahawalpur | 35.7 | 17.6 | 59.4 | 0.0 | 36.0 | 21.6 | 58.9 | 0.0 |
| Bahawalnagar | 36.7 | 22.2 | 66.4 | 0.0 | 32.9 | 22.5 | 67.8 | 0.0 |
| R.Y.Khan | 38.0 | 24.2 | 65.2 | 0.0 | 37.0 | 20.2 | 41.6 | 0.0 |
| D.G. Khan | 32.1 | 24.1 | 61.8 | 0.0 | 37.4 | 26.7 | 69.6 | 0.0 |
| Muzaffar Garh | 36.9 | 23.9 | 0.0 | 0.0 | 34.2 | 22.1 | 68.2 | 0.0 |
| Rajanpur | 37.5 | 25.0 | 60.1 | 0.0 | 36.3 | 28.8 | 53.7 | 0.0 |
| Layyah | 35.0 | 18.0 | 50.0 | 0.0 | 34.0 | 21.0 | 74.0 | 0.0 |
| AVERAGE | 35.44 | 22.57 | 53.95 | 0 | 35.13 | 22.40 | 63.50 | 0 |

Weather forecast for next 7 day in cotton zone

| Division | Dated | 16/10 | 17/10 | 18/10 | 19/10 | 20/10 | 21/10 | 22/10 |
|-------------------|------------|-------|-------|-------|-------|-------|-------|-------|
| | Max.Temp. | 31 | 27 | 29 | 31 | 33 | 34 | 34 |
| Multan | Min.Temp. | 22 | 19 | 19 | 18 | 19 | 20 | 22 |
| | Humidity % | 53 | 55 | 44 | 40 | 35 | 29 | 27 |
| | Max.Temp. | 28 | 28 | 27 | 29 | 32 | 34 | 36 |
| Sahiwal | Min.Temp. | 19 | 18 | 16 | 15 | 16 | 17 | 22 |
| | Humidity % | 65 | 63 | 53 | 44 | 32 | 28 | 23 |
| | Max.Temp. | 34 | 28 | 38 | 32 | 34 | 36 | 37 |
| Bahawalpur | Min.Temp. | 23 | 19 | 18 | 18 | 19 | 20 | 21 |
| | Humidity % | 35 | 56 | 42 | 34 | 33 | 24 | 21 |
| D.C | Max.Temp. | 31 | 28 | 30 | 32 | 33 | 36 | 35 |
| D.G. Khan | Min.Temp. | 22 | 19 | 19 | 19 | 20 | 21 | 22 |
| Kilali | Humidity % | 40 | 44 | 39 | 36 | 33 | 26 | 24 |
| Average Cotton | Max.Temp. | | | | 27 | | | |
| | Min.Temp. | | | | 19.36 | | | |
| Region | Humidity % | | | | 38.5 | | | |

5

Source: timeanddate.com

Summary of weather forecast

Overall weather for seven days in cotton zone during next week will remain

partly cloudy, Sunny & Shiny.

Forecast of Sucking Pests:

Whitefly: This pest flourishes best in hot dry climate with optimum temperature 35-

42 °C with relative humidity below 50%. Maximum lethal temperature for this pest is

45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.44

°C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max.

27.0°C, Min. 19.36°C with humidity 38.5%. Keeping in view above all, it is predicted

that population of this pest will sustain during the coming week.

Jassid: This pest flourishes best in hot and humid climate with optimum temperature

35-40 °C with relative humidity above 70%. Maximum lethal temperature for this

pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max.

35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp.

Max. 27.0°C, Min. 19.36°C with humidity 38.5%. Keeping in view above all, it is

predicted that population of this pest will sustain during the coming week.

Thrips: This pest flourishes best in hot dry climate with optimum temperature 35-40

°C with relative humidity below 60%. Maximum lethal temperature for this pest is 45-

51 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C,

Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max.

27.0°C, Min. 19.36°C with humidity 38.5%. In view above all, it is predicted that

population of this pest will sustain during the coming week.

Mealy Bug: This pest flourishes best in moderate climate with optimum temperature

35-40 °C with relative humidity above 60-80%. Maximum lethal temperature for this

pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with humidity 38.5%. Keeping in view above all, it is predicted that population of this pest will sustain as such on cotton crop during the coming week.


Mites: This pest flourishes best in dry climate with optimum temperature 28-40 °C with relative humidity above 40-50%. Maximum lethal temperature for this pest is 45 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with humidity 38.5%. Keeping in view above all, it is predicted that population of this pest will sustain as such during the coming week.

Dusky Cotton Bug: This pest flourishes best in moderate climate with optimum temperature 28-40 °C with relative humidity above 50-80%. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with humidity 38.5%. In view above all, it is predicted that population of this pest will increase during the coming week.

Pink Bollworm: This pest flourishes best at moderate climate with optimum temperature 32-40 °C with relative humidity above 60-80%. Maximum lethal temperature for this pest is 45-53 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with humidity 38.5%. It is predicted that population of this pest will sustain on cotton during the coming week.

Moth Catches in Pheromone Traps during 2nd Week of October, 2023

| District | Traps Installed | Moth catches/Trap/week | Av. Moth catches/Trap/night |
|--------------|--------------------|------------------------|-----------------------------|
| Multan | 6 | 113 | 2.69 |
| Khanewal | 8 | 158 | 2.82 |
| Vehari | 6 | 111 | 2.64 |
| Lodhran | 6 | 133 | 3.17 |
| Sahiwal | 4 | 18 | 0.64 |
| Pakpattan | 4 | 14 | 0.50 |
| Okara | 4 | 2 | 0.07 |
| Bahawalpur | 8 | 109 | 1.95 |
| Bahawalnagar | 10 | 117 | 1.67 |
| R.Y.Khan | 8 | 101 | 1.80 |
| D.G.Khan | 6 | 67 | 1.60 |
| Rajanpur | 6 | 60 | 1.43 |
| M.Garh | 8 | 62 | 1.11 |
| Layyah | 6 | 55 | 1.31 |
| Bhakkar | 8 | 41 | 0.73 |
| Mianwali | 6 | 39 | 0.93 |
| Faisalabad | 6 | 18 | 0.43 |
| TT. Singh | 6 | 32 | 0.76 |
| Jhang | 4 | 15 | 0.54 |



American Bollworm: This pest flourishes best at dry & moderate climate with optimum temperature 32-40 °C with relative humidity above 50-70%. Maximum lethal temperature for this pest is 45-53 °C. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with humidity 38.5%. It

is predicted that population of this pest will remain sustain on cotton crop during the coming week.

Moth catches on light traps during 2nd week of October, 2023

| Moth Catches on Light Traps/Trap/night During 2nd Week of October, 2023 | | | | | | | | | |
|---|-----------------|-------|-------|----------|--|--|--|--|--|
| District | Traps Installed | A.B.W | S.B.W | Armyworm | | | | | |
| Multan | 3 | 1.0 | 0.0 | 3.0 | | | | | |
| Khanewal | 4 | 1.4 | 0.0 | 5.4 | | | | | |
| Vehari | 3 | 0.0 | 0.0 | 0.9 | | | | | |
| Lodhran | 3 | 0.9 | 0.0 | 2.1 | | | | | |
| Sahiwal | 2 | 0.3 | 0.0 | 0.6 | | | | | |
| Pakpattan | 2 | 0.3 | 0.0 | 0.4 | | | | | |
| Okara | 3 | 0.4 | 0.0 | 0.7 | | | | | |
| Bahawalpur | 4 | 1.6 | 0.9 | 2.7 | | | | | |
| Bahawalnagar | 5 | 1.1 | 0.7 | 3.0 | | | | | |
| R.Y.Khan | 4 | 0.9 | 0.6 | 1.4 | | | | | |
| D.G.Khan | 3 | 0.9 | 0.0 | 1.3 | | | | | |
| M.Garh | 4 | 0.0 | 0.0 | 0.3 | | | | | |
| Rajan Pur | 3 | 0.3 | 0.1 | 1.0 | | | | | |
| Layyah | 3 | 0.0 | 0.0 | 0.0 | | | | | |

Note. 3 moth catches of a pest on 3 consecutive nights in a trap is considered as ETL

Spotted Bollworm: This pest flourishes best at mild & humid climate with optimum temperature 30-40 °C with relative. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with humidity 38.5%. Due to cultivation of more BT varieties, it is predicted that population of this pest will remain as such on cotton crop in the coming week.

Armyworm: This pest flourishes best in mild & humid climate with optimum temperature 30-40 °C with relative humidity below 70-100%. The current weather conditions on overall Punjab basis (temp. Max. 35.44 °C, Min. 22.57°C with R. humidity 53.95%) & forecast for the next week temp. Max. 27.0°C, Min. 19.36°C with

humidity 38.5%. Keeping in view above all, it is predicted that population of this pest will sustain during the coming week.

CLCuV: The incidence of this disease is expected to sustain as such on cotton crop during the next week.

RECOMMENDATIONS

WHITEFLY MANAGEMENT

- 1. Conduct regular pest scouting of Whitefly.
- 2. Control the pest without delay when population reaches ETL i.e. nymphs or adults or both 5 per leaf.
- 3. Control weeds in and outside the field especially near water channels and fallow lands.

Alternate host plants of whitefly

| Crops | Vegetables | Orchards | Weeds/ | Trees |
|-----------|-------------|-------------------------|-------------|---------|
| | | | Ornamentals | |
| Sunflower | Okra | Citrus | Gardenia | Shisham |
| Tobacco | Brinjal | Litchi | Mako | Shareen |
| | Cucurbits | Pomegranate | Maina | |
| | Tomato | Ber (<i>Zizyphus</i>) | Karund | |
| | Cabbage | Guava | Lehli | |
| | Cauliflower | Mulberry | | |
| | Peas | Papaya | | |
| | Potato | | | |
| | Onion | | | |
| | Spinach | | | |

JASSID MANAGEMENT

- 1. Eradicate weeds.
- 2. Conduct Pest Scouting at least twice a week.
- 3. Use *Chrysoperla* Cards @ 80-90 cards per acre having 20-25 eggs on each card which are available at AARI Faisalabad, Sahiwal, Okara and Vehari. (Repeat this process after every two weeks).

THRIPS MANAGEMENT

- 1. Conduct Pest Scouting regularly at least twice a week.
- 2. Avoid intercropping of cotton crop with vegetables.
- 3. Control weeds in and outside the field.

MEALY BUG MANAGEMENT

- Visit the fields daily to detect the infestation for effective management of mealy bug well in time.
- 2. Keep field and water courses etc. free from weeds especially Itsit & Hazaardani, kangi booti, aksin etc.
- To avoid its shifting from one place to another, keep the horticultural nurseries free from its infestation. Spray such infested nurseries before shifting the plants.
- 4. Avoid water stress to the plants.
- 5. Avoid repeated visits of workers from infested fields to healthy fields.
- 6. Apply the localized chemical spray on infected along with its 2 meter surrounding area.

Alternate host plants of cotton mealy bug

| Crops | Vegetables | Ornamentals | Weeds | Orchards |
|-----------|------------|-------------|-------------------|----------|
| Sunflower | Okra | China Rose | Hazar Dani Citrus | |
| Tobacco | Brinjal | Huddle | Amarantus | Mulberry |
| Jantar | Tomato | Cotton Rose | Bhakra | Ficus |
| Sesame | Chillies | Gulchain | Mako | Ber |
| | Pumpkin | Lantana | Sueda | |
| | | Din Ka Raja | Itsit | |
| | | Rat Ki Rani | Karund | |
| | | Anthorium | Aksen | |
| | | Gul-e-Daudi | Bathu | |
| | Gainda | | Puth Kanda | |
| _ | | | Kanghi | _ |

MANAGEMENT OF DUSKY COTTON BUG

- 1. Keep the special observation/ pest scouting when the boll opening starts.
- **2.** Use of soft contact poison on the above ETL.

MANAGEMENT OF ARMY WORM

- Keep the fields clean from weeds especially Its it which is a preferred host of armyworm.
- 2. Observe the field on regular basis and under close observation to assess its attack and treat the patch at earliest if necessary.
- 3. Avoid sowing of Jantar near cotton fields.
- 4. Regular pest scouting at least twice a week.
- 5. Hand picking and destruction of egg masses at initial stage as initially its attack starts in patches and is clearly visible.

CLCuV MANAGEMENT

- 1. Use CLCuV tolerant varieties for sowing.
- 2. Keep Cotton fields free from all kinds of weeds.
- 3. Eradicate other alternate host plants of CLCuV and dispose them off carefully.
- 4. Thinning out and destruction of Virus affected plants.
- 5. Remove CLCuV affected plants in less than 60 DAS (Days after sowing) crop.
- 6. Hoeing after each irrigation or rain at wattar condition till canopies permit.
- Apply post emergence Glyphosate for both broad and narrow leaved weeds and Glyphosate for grassy weeds with shield and use Flat Fan Nozzle, if needed.
- 8. Irrigate the fields when needed keeping in view weather conditions e.g. temperature, rainfall, soil water holding capacity and plant need.

- 9. Balanced use of Fertilizer i.e. N.P.K. along with foliar growth promoters if necessary.
- 10. Keep transmission vector (whitefly) at the low ebb

Alternate host plants of CLCuV vector (whitefly)

| Crops | Vegetables | Ornamentals | Weeds |
|-----------|------------|-------------|--------------|
| Sunflower | Okra | Gurhal | Leh |
| Melon | Brinjal | Chambeli | Lehli |
| Tobacco | Chillies | | Mako |
| | Tomato | | Maina |
| | Potato | | Karund/Bathu |
| | Cucumber | | Gardenia |
| | | | Hazardani |
| | | | Rattanjot |
| | | | Sun Kukra |

Economic threshold levels of cotton pests

| Insect pests | Economic threshold levels | |
|--------------------|---|--|
| Jassid | 1 Adult or Nymph per leaf | |
| Whitefly | 5 Adults or Nymphs or both per Leaf | |
| Thrips | 8-10 Adults or Nymphs per Leaf | |
| Mealy bug | 0 (on appearance) | |
| Mites | On damage appearance | |
| Spotted Boll worm | 3 Larvae/ 25 plants. | |
| Pink Boll worm | 5 Larvae /100 bolls. | |
| American boll worm | 5 brown eggs or 3 small larvae or both 5 per 25 plants on non Bt varieties. 2 larvae of 2 nd instar per 25 plants on Bt varieties. | |
| Army worm | 0 (On appearance) | |
| Aphid | Spray on visible damage on top terminals. | |
| Dusky Cotton bug | 10 per leaf/bud | |